

# Wiener Polynomials of some Chemically Interesting Graphs

Bo-Yin Yang

Yeong-Nan Yeh

Department of Mathematics

Institute of Mathematics

Tamkang University, Tamsui,

Academia Sinica\*, Nankang

Taipei County, Taiwan, R.O.C.

Taipei, Taiwan, R.O.C.

March 10, 2003

## Abstract

The chemist Harold Wiener found  $\mathcal{W}(G)$ , *the sum of distances between all pairs of vertices* in a connected graph  $G$ , to be useful as a predictor of certain physical and chemical properties. The  $q$ -analog of  $\mathcal{W}$ , called the *Wiener Polynomial*  $\mathcal{W}(G; q)$ , is also useful but has few existing useful formulas. We will evaluate  $\mathcal{W}(G; q)$  for certain graphs  $G$  of chemical interest.

Keywords: Wiener Polynomial, polygonal chains, hexagonal carpets

AMS Classification: Combinatorics (05)

---

\*Partially supported by National Science Council grant NSC89-2115-M-001-024.

# 1 Introduction

A structural formula in organic chemistry corresponds naturally to a connected graph, each non-hydrogen atom to one vertex. Distance between vertices corresponds to the number of bonds between two atoms, which quantum-mechanically influences physical and chemical properties. In 1947, Harold Wiener proposed a formula ([30]) for boiling points of alkanes, after some heuristic reasoning:

$$\text{b.p.} = \alpha\mathcal{W} + \beta w_3 + \gamma \quad (1)$$

Where (we quote) “ $\mathcal{W}$  is the sum of distances and  $w_3$  the number of pairs of vertices three apart.” The correlation was surprisingly good, so he published a few more papers on the same topic.  $\mathcal{W}$  was rediscovered in the 1970’s, and in time the mathematical properties were probed to some detail, the most notable result of this period being the Merris-McKay theorem relating Wiener numbers to eigenvalues of trees ([14]). Descriptions of its applications have abound in the literature since, numbering hundreds of articles. “Topological indices” — the chemists’ name for a function on a graph — in general and their applications are discussed in chemical texts such as [1, 18, 28]; surveys on Wiener numbers and their applications etc. can be found in [7, 15, 18, 33]. Other treatments are in [2, 3, 4, 5, 9, 10, 11, 16, 17, 21, 22, 23, 25, 27].

It is natural to consider generating functions (“counting polynomials”)

when studying something in a combinatorial structure in aggregate, a direct extension to Wiener’s idea was mentioned early by Haruo Hosoya, and formalized as well as investigated in some detail by Sagan and Yeh:

**Definition 1** ([11, 24]) *Let  $d_G : V \times V \rightarrow \mathbb{N}$  be the function<sup>1</sup> representing minimal distance between vertices of the connected graph  $G = (V, E)$ . The counting polynomial of distances on  $G$ —the **Wiener Polynomial** of  $G$ —is*

$$\mathcal{W}(G; q) \equiv \sum_{\{u,v\} \in \binom{V}{2}} q^{d_G(u,v)};$$

*we also define  $\mathcal{W}(u|G; q) \equiv \sum_{v \in V} q^{d_G(u,v)}$  and  $\mathcal{W}(S_1, S_2|G; q) \equiv \sum_{u \in S_1, v \in S_2} q^{d(u,v)}$ .*

*These are  $q$ -analogs of  $\mathcal{W}(G)$ ,  $\mathcal{W}(u|G)$ , and  $\mathcal{W}(S_1, S_2|G; q)$  ([13]). Each*

*quantity relate to its  $q$ -analog in the usual manner, e.g.  $\mathcal{W}(G) = \frac{d}{dq} \mathcal{W}(G; q)|_{q=1}$ .*

Like most generating functions, Wiener polynomials has some independent interest: an example ([24]) is its linking the absolute Poincaré polynomials of a Coxeter group and its induced graph. Knowing how to compute Wiener polynomials would also yield Wiener’s “polarity number”  $w_3$ . Wiener number laid dormant for a decade and a half before being rediscovered at least partly due to the scarcity of efficient formulas for  $w_3$  in chemically interesting graphs. There are a few other practical situation in which points at a given distance needs to be counted.

---

<sup>1</sup> *We shall write  $\mathbb{N}$  for the set of non-negative integers and  $\mathbb{P}$  positive ones in this paper; other notations for convenience we will use are  $u$  for the singleton set  $\{u\}$ , and  $d$  for  $d_G$  when confusion is unlikely!*

Wiener polynomials of simple graphs are well-known:

$$\mathcal{W}(C_{2n}; q) = n \left[ \frac{(q+1)(q^n-1)}{q-1} - 1 \right]; \quad (2)$$

$$\mathcal{W}(C_{2n+1}; q) = (2n+1)q \frac{q^n-1}{q-1}; \quad (3)$$

$$\mathcal{W}(P_n; q) = \sum_{j=1}^{n-1} (n-j)q^j = \frac{q(q^n-1)}{(q-1)^2} - \frac{nq}{(q-1)}. \quad (4)$$

Sagan and Yeh ([24]) discussed certain relationships that allows us to construct Wiener polynomials of graphs formed by some binary operations from simpler graphs, such as the paths  $P_n$  and cycles  $C_n$ . The most useful of these relations pertain to the Cartesian product:

$$\mathcal{W}(G \times H; q) = 2\mathcal{W}(G; q)\mathcal{W}(H; q) + |V(H)|\mathcal{W}(G; q) + |V(G)|\mathcal{W}(H; q).$$

In particular for the  $m \times n$  Chessboard  $\text{Cb}_{m,n} \equiv P_m \times P_n$ :

$$\mathcal{W}(\text{Cb}_{m,n}; q) = \frac{[2q(q^m-1) - m(q^2-1)][2q(q^n-1) - n(q^2-1)]}{2(q-1)^4} - \frac{mn}{2}. \quad (5)$$

The authors ([12, 13, 31, 33]) have presented ways to compute Wiener numbers (and occasionally polynomials) for some chemically useful graphs. However the computation of Wiener polynomials often present taller obstacles than when dealing with Wiener numbers, where neat formulas often result by cancellation. When manipulating generating functions this can become anywhere between difficult to impossible. For example, this elegant

result is not easily generalizable:

**Proposition 1 (Merris ([14]))** *Let  $A(T)$  be the adjacency matrix of the tree  $T$  and  $D(T)$  be the diagonal matrix of the same size with the degree as the entry corresponding to each vertex. The Laplacian matrix  $L(T)$  is defined as  $D(T) - A(T)$ . Merris showed that*

$$\mathcal{W}(T) = |V(T)| \sum \frac{1}{\lambda},$$

where  $\lambda$  ranges over all eigenvalues of  $L(T)$ .

In Sec. 2 we will treat the problem of how to compute Wiener Polynomials of polygonal chains; these graphs (“motley chains”) are the abstractions of aromatic compounds and their Wiener numbers and polynomials are of significant interest. In Sec. 3 we will deal with Wiener polynomials for some regular 2-dimensional hexagonal patterns. These graphs (“hex carpets”) depicts slices of graphite.

## 2 Wiener Polynomials of Polygonal Chains

**Definition 2 ([32])** *A motley chain is a graph of concatenated, or edge-sharing, polygons. Given  $n$  ordered pairs of non-negative integers  $S = (a_1, b_1), \dots, (a_n, b_n)$ , we may create a graph as follows: take the graph  $\text{Cb}_{2, n+1} = P_2 \times P_{n+1}$  and subdivide the upper and lower edges by inserting  $a_j$  and  $b_j$  extra vertices. respectively (see Fig. 1). The resulting graph is the motley*


chain associated with  $S$ . Since only  $a_1 + b_1$  and  $a_n + b_n$  matters we will denote equivalence classes as:

$$S = \left\langle \begin{matrix} & a_2 & a_3 & \cdots & a_{n-1} & & \\ (a_1 + b_1) & & & & & & (a_n + b_n) \\ & b_2 & b_3 & \cdots & b_{n-1} & & \end{matrix} \right\rangle.$$


When each cycles is of even order (as in Figure 1(b)), we call it an **even motley chain**. Let integers  $k_i$  and  $j_i$  satisfy  $|j_i| < k_i$ ,  $i = 1 \dots n$ , then we write

$$E = k_0(k_1)_{j_1}(k_2)_{j_2} \cdots (k_n)_{j_n}k_{n+1}$$


as a representation for the even motley chain in which the  $i$ -th polygon has  $2(k_i + 1)$  vertices, of which  $k_j + j_i + 1$  are along the lower edge, or  $2j_i$  more than those along the upper edge. For obvious reasons,  $k_i$  is called the length of the  $i$ -th cell.  $\mathcal{K} \equiv \sum_{j=0}^{n+1} k_j$  is the **total length** of the chain.



(a) A Motley Chain



(b) An Even Motley Chain



(c) Another rendering of (b), depicting a real compound

Figure 1: “Chains of motley gems”

It is useful to compare a “straight”  $E_0 = k_0 k_1 \cdots k_{n+1}$  even motley chain (all the  $j_i$ ’s are 0) to  $P_{\mathcal{K}+1} \times P_2$  (a chain of  $\mathcal{K}$  squares). Noting that when




Figure 2: Evaluating Wiener Polynomials of straight even motley chains

any  $2k + 2$ -gon is cut into  $k$  squares, distances between other vertices do not change, we get (using Eqs. 2 and 5):

**Proposition 2 ([33])** *The Wiener polynomial of the straight even motley chain  $E_0$  to be (independent of the order of the polygons):*

$$\mathcal{W}(E_0; q) = \mathcal{W}(\text{Cb}_{\mathcal{K}+1,2}) + \sum_j \frac{q+1}{q-1} \left[ (k_j - 1)q \frac{q^{k_j+1} - 1}{q-1} - (k_j + 1)q^2 \frac{q^{k_j-1} - 1}{q-1} \right]. \quad (6)$$

The general idea will be to start with  $E_0 = k_0 k_1 k_2 \cdots k_n k_{n+1}$ , which is  $E = k_0 (k_1)_{j_1} (k_2)_{j_2} \cdots (k_n)_{j_n} k_{n+1}$  with every kink straightened out, and morph it into  $E$  one rotation at a time, starting from the left: the  $\ell$ -th step in the process is rotating the  $i_\ell$ -th polygon from being “straight” into the “bent” position specified by  $(k_{i_\ell})_{j_{i_\ell}}$ . But first we need this generally useful result:

**Proposition 3 (Shelling Lemma, [33])** *Let  $G = (V, E)$  be a connected graph, and partition its vertex set  $V$  into  $V_0 \uplus V_1 \uplus V_2 \uplus \cdots \uplus V_k$  in such a way that the restriction of  $d_G \equiv d$  to  $G_j \equiv G|_{V_j}$  is the same as  $d_{G_j}$  (hence,*

each  $V_j$  is connected). Also let  $\mathcal{G}_j$  be the subgraph induced by  $\bigcup_{i=1}^j V_i$ , then

$$\begin{aligned} \mathcal{W}(G; q) &= \sum_{j=0}^k \mathcal{W}(G_j; q) + \sum_{0 \leq i < j \leq K} \mathcal{W}(G_i, G_j; q) \\ &= \mathcal{W}(G_0; q) + \sum_{j=1}^k \left[ \sum_{u \in V_j} \mathcal{W}(u | \mathcal{G}_j; q) - \mathcal{W}(G_j; q) - |V_j| \right]. \quad (7) \end{aligned}$$




Figure 3: Changes after one rotation

The portion of the chain to the left of a bend will be the “residue”  $R$ , and the remaining straight portion on the right the “tail”  $T$ . Now introduce just one bend (nonzero  $j$ ) into a straight chain, so that  $R$  is also straight.

**Lemma 4** *Let there be only the one bend  $(k_i)_{j_i}$  (see Fig. 3), where  $i = i_1$  is the only index corresponding to nonzero  $j_i$ . Then*

$$\mathcal{W}(E_0; q) - \mathcal{W}(E; q) = q^{k_i+2-|j_i|} (q+1) (2q^{|j_i|} - q - 1) \frac{q^{K'} - 1}{q-1} \frac{q^K - 1}{q-1},$$

where  $K$  and  $K'$  are the length of  $R$  and  $T$ , respectively.

**Proof:** Use Prop. 3 with  $R$ , the polygon of the bend, and  $T$  as the parts.

When we rotate the  $i$ -th polygon only the distances from  $R$  to  $T$  change.

These distances originally contribute to the Wiener polynomial the sum

$$2(q+1)q^{k_i+2} \left( \frac{q^K - 1}{q - 1} \right) \left( \frac{q^{K'} - 1}{q - 1} \right)$$

which now becomes (all pertinent distances going through the marked path):

$$(q+1)^2 q^{k_i+2-|j_i|} \left( \frac{q^K - 1}{q - 1} \right) \left( \frac{q^{K'} - 1}{q - 1} \right).$$

□

Before we proceed, we need to tidy up some notations:

$$\begin{aligned} K_\ell &\equiv \sum_{h=0}^{i_\ell-1} k_h, \\ K'_\ell &\equiv \sum_{h=i_\ell+1}^{n+1} k_h, \\ \hat{j}_\ell &\equiv \begin{cases} |j_{i_\ell}|, & \text{if } j_{i_\ell} j_{i_\ell-1} > 0, \\ |j_{i_\ell}| - 1, & \text{if } j_{i_\ell} j_{i_\ell-1} < 0 \\ 0 & \text{else.} \end{cases} \\ L_\ell &\equiv K_\ell - \sum_{h=0}^{\ell-1} (\hat{j}_h) \end{aligned}$$

As we shall see,  $K_\ell$  and  $K'_\ell$  are the sizes of leading and tail partial chains ( $K_0 \equiv 0$ ).  $\hat{j}_\ell$  represent the “*kink*” at the  $i_\ell$ -th polygon, and  $L_\ell$  is the ‘true’ length of the leading partial chain. Now to account for more than one bend

or kink in a chain (see Fig. 4). For each bend  $(k_{i_\ell})_{j_{i_\ell}}$ , let  $u_\ell$  be the top




Figure 4: The two Segments of  $R$

vertex between the  $(i_\ell - 1)$ -th polygon (and the rest of the remaining chain  $R_\ell$ ) to the  $i_\ell$ -th polygon if  $j_\ell > 0$ , and the bottom vertex if  $j_\ell < 0$ . Also let  $\mathcal{X}_\ell(q) \equiv \mathcal{W}(u_\ell | R_\ell; q)$ , which we will need later:

**Lemma 5**

$$\begin{aligned}
& \mathcal{W}(E_0; q) - \mathcal{W}(E; q) \\
&= \sum_\ell (q+1) \left( \frac{q^{K'_\ell} - 1}{q-1} \right) \left\{ \left( \frac{q^{K_\ell - K_{\ell-1}} - 1}{q-1} \right) q^{k_{i_\ell} + 2 - |j_{i_\ell}|} (2q^{|j_{i_\ell}|} - q - 1) \right. \\
&\quad \left. + (q^{K_\ell - K_{\ell-1}} - q^{K_\ell - K_{\ell-1} - \hat{j}_{i_\ell}}) \frac{\mathcal{X}_{\ell-1}(q)}{q} \right\} \tag{8}
\end{aligned}$$

**Proof:** In general the residue  $R_\ell$  bends, but comprises one straight segment (effectively two paths of length  $K_\ell - K_{\ell-1}$ ) attached to the  $\ell$ -th bend, and a remainder  $R_\ell - 1$  beyond it. Use the Shelling Lemma with the polygons at the kinks and the straight segments in between as the parts. So the  $\ell$ -th

rotation decrease the Wiener polynomial by

$$q^{k_{i_\ell}+2-|j_{i_\ell}|}(q+1)(2q^{|j_{i_\ell}|-q-1})\frac{q^{K'_\ell}-1}{q-1}\frac{q^{K_\ell-K_{\ell-1}}-1}{q-1}+(\text{contribution from } R_{\ell-1}).$$

The term involving  $R_{\ell-1}$  can be deduced thus: any shortest path from  $T$  into  $R_{\ell-1}$  must traverse the path  $\overline{uv}$ . So the partial Wiener sum starts out as (again using the Shelling Lemma)

$$\mathcal{W}(u_{\ell-1}|R_{\ell-1}; q) \cdot q^{K_\ell-K_{\ell-1}-|j_{i_{\ell-1}}|+k_{i_\ell}+1} \cdot (q+1)\frac{q^{K'_\ell}-1}{q-1},$$

and is scaled down by a factor of  $q^{\hat{j}_{i_\ell}}$ , where  $\hat{j}$  is the *kink* defined earlier.  $\square$


Now we are ready for the finale:

**Theorem 6** *Let  $E = k_0(k_1)_{j_1}(k_2)_{j_2} \cdots (k_n)_{j_n}k_{n+1}$  be an even motley chain, with exactly  $j_{i_1}, j_{i_2}, \dots, j_{i_m}$  non-zero among all the  $j_i$ . With notations as above, the Wiener polynomial of  $E$  is given by:*


$$\begin{aligned} \mathcal{W}(E; q) &= \mathcal{W}(\text{Cb}_{\mathcal{K}+1,2}; q) + \sum_j \frac{q+1}{q-1} \left[ (k_j-1)q\frac{q^{k_j+1}-1}{q-1} - (k_j+1)q^2\frac{q^{k_j-1}-1}{q-1} \right] \\ &- \sum_\ell (q+1) \left( \frac{q^{K'_\ell}-1}{q-1} \right) \left\{ \left( \frac{q^{K_\ell-K_{\ell-1}}-1}{q-1} \right) q^{k_{i_\ell}+2-|j_{i_\ell}|} (2q^{|j_{i_\ell}|-q-1}) \right. \\ &+ \left. (q+1) (q^{K_\ell-K_{\ell-1}} - q^{K_\ell-K_{\ell-1}-\hat{j}_\ell}) \left[ \frac{q^{L_{\ell-1}}-1}{q-1} + \sum_{h=1}^{\ell-2} q^{L_{\ell-1}-L_h} \left( \frac{q^{\hat{j}_h}-1}{q-1} \right) \right] \right\} \end{aligned} \quad (9)$$

From Eq. 9 we can deduce all previous results about Wiener numbers and polynomials of an even motley chain; actually Eq. 8 suffices when we note


that there are  $2K_\ell$  vertices in  $R_\ell$  and hence that many terms in  $\mathcal{X}_\ell(q)$ .



(a) A straight Chain:  $\mathcal{X} = q(q+1) \left( \frac{q^{18} - 1}{q - 1} \right)$



(b) Curving one way:  $\mathcal{X} = q(q+1) \left( \frac{q^{17} - 1}{q - 1} + q^{15} \right)$



(c) Curving the other way:  $\mathcal{X} = q(q+1) \left( \frac{q^{16} - 1}{q - 1} + q^{14} \frac{q^2 - 1}{q - 1} \right)$

Figure 5: Partial Wiener Polynomials for almost-straight Even Chains

**Proof:** We still need to find  $\mathcal{X}_i(q)$ . First we look at a few diagrams: As for hex chains, it's easy to see that  $\mathcal{X}_i^*(q) = q(q+1) \left( \frac{q^{K_i} - 1}{q - 1} \right)$ . When the chain develop a kink, the chain effectively became shorter. Note that the upper and lower edges differ, due to the starting location  $u$  (which is on top or bottom when  $j_i$  is positive or negative, respectively).  $\mathcal{X}$  becomes

$q(q+1) \left( \frac{q^{K_i-\hat{j}} - 1}{q-1} + q^p \frac{q^{\hat{j}} - 1}{q-1} \right)$  after one kink is introduced, where  $p$  is the ‘effective length’ between the kink and the endpoint  $u$ , and  $\hat{j}$  is the amount of the kink, i.e.  $|j|$  when it is the first curve of two in the same direction, and  $|j| - 1$  otherwise. Fig. 6 shows what kind of terms appears as the polygons are rotated one by one to create our motley chain. We can now verify the important relation:

$$\mathcal{X}_\ell(q) = q(q+1) \left[ \frac{q^{L_\ell} - 1}{q-1} + \sum_{h=1}^{\ell-1} q^{L_\ell - L_h} \left( \frac{q^{\hat{j}_{i_h}} - 1}{q-1} \right) \right]. \quad (10)$$




Figure 6: A Chain with many Turns:  $\mathcal{X} = q(q+1) \left( \frac{q^{14} - 1}{q-1} + q^{12} \frac{q^2 - 1}{q-1} + q^8 + q^3 \right)$

Now sum up the terms in straightforward fashion to get Eq. 9.  $\square$

We sketch with some examples how to obtain the Wiener polynomials of polygonal chains involving odd cycles. Following [32], we term a motley chain *zigzagging* or *straight* (Fig. 7) if for at least one representation  $S = (a_1, b_1), \dots, (a_n, b_n)$ , we have

$$\left\{ \sum_{i=1}^j (a_i - b_j) \mid 1 \leq j \leq n \right\} \subset \{0, 1\} \text{ or } \{0, -1\}.$$




Figure 7: A ‘Straight’ Motley Chain

We aim to work out the Wiener Polynomials of motley chains, starting with straight ones and working our way down the line, adding terms for each bend or twist in a similar way to the above.


**Lemma 7 (Subdivision of even cycle)** *A polygonal chain of one  $(2m + 1)$ -cycle, one  $2(k+1)$ -cycle, and one  $(2n+1)$ -cycle has its Wiener polynomial given by*

$$\begin{aligned} & \mathcal{W} \left( \left\langle \begin{matrix} (2m-3) & k-1 & (2n-3) \\ & k-1 & \end{matrix} \right\rangle; q \right) \\ &= (2n+1)q \frac{q^n - 1}{q-1} + (k+1) \left[ \frac{(q+1)(q^n - 1)}{q-1} - 1 \right] + (2m+1)q \frac{q^m - 1}{q-1} - 2q \\ &+ \frac{q^{n+1+m+k} + 4q^{n+m+k} - q^{n+m+k-1} - 2q^{2+k} - 2q^{1+k} - 4q^{2+m} + 4q^3 + 4q^2 - 4q^{n+2}}{(q-1)^2} \end{aligned}$$


*In particular the Wiener polynomial of fused  $(2n+1)$ - and  $(2m+1)$ -cycles, or  $\mathcal{W}(\langle(2m-3), (2n-3)\rangle; q)$  is given by*

$$\mathcal{W}(C_{2n+1}; q) + \mathcal{W}(C_{2m+1}; q) - q + \frac{q^{m+n}(q^2 - 1) + 4q^3(q^{m-1} - 1)(q^{n-1} - 1) - 2q^3 + 2q^2}{(q-1)^2}.$$

Denote a straight chain of  $n$  repetitions of fused cycles of  $n_1, n_2, \dots, n_k$  sides in that order to be  $\mathcal{Z}_{n_1, n_2, \dots, n_k}(n)$ . So a straight motley chain com-



(a) Fragments



(b) Process of subdivision

Figure 8: Septagon-Pentagon chains

posed of  $n$  pairs of alternating pentagons and septagons would be  $\mathcal{Z}_{5,7}(n)$ . Since the chain fragment consisting of one pentagon and one septagon can be created from partitioning a decagon (which has *length* 4), and the difference between its Wiener polynomial and 4 fused squares is  $\mathcal{W}(\langle 1, 3 \rangle; q) - \mathcal{W}(\text{Cb}_{5,2}; q) = -q^5 + 3q^3 - 2q$ , One would expect  $\mathcal{W}(\mathcal{Z}_{5,7}(n); q)$  to be given by  $\mathcal{W}(\text{Cb}_{10n+1,2}; q) - nq(q^2 - 1)(q^2 - 2)$ , but this is not so. As we subdivide one of the decagons we have the four blobbed vertices marked in Fig. 2 above, each getting closer to each of the circled ones by 1.

So, to get  $\mathcal{W}(\mathcal{Z}_{2k+1,2m+1}(n); q)$ , we have to add in addition to the differences between the fragments, which is  $n [\mathcal{W}(\langle 2k - 3, 2m - 3 \rangle; q) - \mathcal{W}(\text{Cb}_{k+m,2}; q)]$ ,

the term

$$\sum_{j=1}^{n-1} \left( \frac{q^k - 1}{q - 1} \frac{q^{(k+m-1)j} - 1}{q - 1} + \frac{q^{m-1} - 1}{q - 1} \frac{q^{(k+m-1)j-k+1} - 1}{q - 1} \right),$$

which ends in this lemma, which we can use to obtain the Wiener Polynomial of all generic motley chains inductively.

**Lemma 8 (one zigzagging straight chain)**

$$\begin{aligned} & \mathcal{W}(\mathcal{Z}_{2k+1,2m+1}(n); q) \\ = & \mathcal{W}(\text{Cb}_{n(k+m-1)+1,2}; q) + n[\mathcal{W}(\langle 2k - 3, 2m - 3 \rangle; q) - \mathcal{W}(\text{Cb}_{k+m,2}; q)] \\ - & \frac{n-1}{(q-1)^2} (q^{m-1} + q^k - 2) + \frac{(q^k - 1)q^{k+m-1} + (q^{m-1} - 1)q^m}{(q-1)^3} (q^{(k+m-1)(n-1)} - 1) \end{aligned}$$

### 3 2-dimensional Patterns

Mathematical chemists posed the question of computing the Wiener numbers of graphs with 2-dimensional hexagonal patterns (called *hexagonal animals* or *hex carpets* – they resemble a hexagon tiling of the floor) a long time ago. Eventually a solution was found via this idea:

**Definition 3 (Squaring)** *A graph  $G = (V, E)$  representing a given hexagonal animal is said to be **embedded** in the set of lattice points  $\mathbb{Z}^2$  if  $V \subset \mathbb{Z}^2$ ,*

$$E \subset \mathcal{L} \equiv (\{(i, j), (i + 1, j)\} | i, j \in \mathbb{Z}\} \cup \{(i, j), (i, j + 1)\} | i, j \in \mathbb{Z}\}$$

(i.e. , and all edges in  $E$  can be drawn as line segments of length 1).

A graph  $G$  embedded in the lattice points is said to be **partitioned into ‘row-paths’**  $R_j$  if  $V = \bigsqcup R_j$  such that each  $R_j$  contains all vertices with ordinate  $j$  and is isomorphic to a path. It is easy to see that any lattice embedding of a hex carpet with a partition into row-paths will show each 6-cycle (hex) as the boundary of a domino (horizontal  $2 \times 1$  rectangle). Let  $G = (V, E)$  be a hexagonal animal lattice-embedded, then the subgraph of the lattice grid induced by  $V$

$$G^\square \equiv (V, \binom{V}{2} \cap \mathcal{L}) = (\mathbb{Z}^2, \mathcal{L})|_V,$$

is defined as the **squaring** of  $G$ .

We will in the rest of the section demonstrate how to compute the Wiener polynomial for one case of the hex carpets.




Figure 9: A Rectangular Carpet

One basic families of hex carpets is shown in Fig. 9 with its squaring.

**Lemma 9 (Change of distance induced by squaring)** *If  $u = (0, 0) \in R_0$  and  $v = (i, j) \in R_j$  are two vertices  $j(> 0)$  rows apart in a lattice embedding of  $G = (V, E)$  and the row-path partition  $V = R_0 \uplus R_1 \uplus \dots \uplus R_k$ , then  $d_{G^\square}(u, v) = |i| + j$ ; and*

$$d|_G(u, v) = \begin{cases} |i| + j & j \leq |i|; \\ |i| + j + 2 \left\lfloor \frac{j - |i| + 1}{2} \right\rfloor & j > |i|, \{(0, 0), (0, 1)\} \notin E; \\ |i| + j + 2 \left\lfloor \frac{j - |i|}{2} \right\rfloor & j > |i|, \{(0, 0), (0, 1)\} \in E. \end{cases} \quad (11)$$

We present Fig. 10 in lieu of a formal proof:




Figure 10: Effects of Squaring on distances in carpets

Wiener polynomials of chessboards are easy. The following will show that careful manipulations of the changes induced by during squaring will yield the Wiener polynomial of any hex carpet in each of the three major families (see [12]), using similar techniques as that for Wiener numbers.

We compute here as an example the Wiener polynomial of the carpet  $R_{n,k}$ . Taking the squaring of  $R_{n,k}$  we get the chessboard  $\text{Cb}_{2n+1,2k}$ , whose Wiener polynomial  $\mathcal{W}(\text{Cb}_{2n+1,2k}; q)$  is given by

$$\frac{[2q(q^{2n+1} - 1) - (2n + 1)(q^2 - 1)][2q(q^{2k} - 1) - 2k(q^2 - 1)]}{2(q - 1)^4} - k(2n + 1).$$

We hit a snag. Instead of differences as when dealing with Wiener numbers, we must find the actual terms of the respective Wiener Polynomials. Cancellation becomes difficult and patterns can be hard to locate.

**Lemma 10** *The difference terms in the Wiener polynomial induced between the marked vertex  $p$  and the next  $\ell$  rows of vertices (as shown in Fig. 10) is given by:*

$$\begin{aligned} \Delta_p^+(\ell; q) &= q^3 \left[ \frac{1 - q^{2\ell}}{(1 - q^2)^2} - \frac{\ell q^{2\ell}}{1 - q^2} \right] + q^4 \left[ \frac{1 - q^{2\ell-2}}{(1 - q^2)^2} - \frac{(\ell - 1)q^{2\ell-2}}{1 - q^2} \right] \quad (\text{'before'}) \\ &\quad - q \left[ \left( \frac{1 - q^\ell}{1 - q} \right)^2 - \frac{1 - q^\ell}{1 - q} + \frac{1 - q^{2\ell}}{1 - q^2} \right] \quad (\text{'after'}) \\ &= -\frac{\ell q^{2\ell+2}}{1 - q} - \frac{q^{2\ell+2}(q + 2)}{(1 + q)(1 - q)^2} + \frac{q^{\ell+1}(1 + q)}{(1 - q)^2} - \frac{q}{(1 + q)(1 - q)^2}, \end{aligned} \quad (12)$$

**Proof:** By direct summation of the patterns in Fig. 10.  $\square$

The formula above works when the vertex in question is on the “far” side of a hexagon; move it up one row (or look the other way), then we get  $\Delta_p^-(\ell; q)$  which is equal to  $q \Delta_p^+(\ell - 1; q)$ .

If we now sum over the whole of  $\hat{R}_{n,k}$ , always taking the pyramids of numbers (powers of  $t$ ) upwards, then we get a differential of

$$\sum_{\ell=1}^{2k-1} \left[ \left( n + \frac{1 + (-1)^\ell}{2} \right) \Delta_p^+(\ell; q) + \left( n + \frac{1 - (-1)^\ell}{2} \right) \Delta_p^-(\ell; q) \right],$$

which is rather hard to work with, and we do better this (equivalent) way:

$$\sum_{\ell=1}^{2k-1} n [\Delta_p^+(\ell; q) + \Delta_p^-(\ell; q)] + \sum_{\ell=1}^{k-1} [\Delta_p^+(2\ell; q)(1+q)].$$

Now we evaluate the sums in  $\ell$  using of Eq. 12 and get:

**Lemma 11** *The total difference in Wiener Polynomials induced between the hex carpet  $R_{n,k}$  and the row-paths which contain it is given by*

$$\begin{aligned} \Delta^* \mathcal{W}(R_{n,k}) &= \frac{nq}{(1-q)^3} \left[ (1-q^{2k})^2 (1+q) - 2k(1-q)(1-q^{4k}) \right] \\ &+ \frac{2q^{4k+2}}{(1-q)^2(1-q^2)} - \frac{kq}{(1-q)^2} - \frac{q^{2k+1}(q+1)}{(1-q)^3} \\ &+ \frac{q^{4k+2}(2q^4 + q^3 + 2q^2 + q + 2) + (q + 2q^3 + 2q^4 + 2q^5 + q^7)}{(1+q)(1-q)^3(1+q^2)^2} \end{aligned} \tag{13}$$

The formula above sums up all the pyramid-patterns of difference terms, assuming them to be complete throughout. Which it isn't, unfortunately, since the hex carpets have finite width. Ergo we must adjust for the incompleteness of these pyramid around the edges. As in [12], we would take two rows, and consider exactly which terms in these patterns 'disappear over the edges'. Let's call the row-paths in  $R_{n,k}$  (in order)  $L_0, M_0, L_1, M_1, \dots, L_{k-1}, M_{k-1}$ .

We look at just one end of the rows  $L_0$  and  $M_m$  (see Fig. 11), and call the difference terms  $X_m$  as in [12].




Figure 11: “Edge Effects”, as seen before and after the squaring

$$\begin{aligned}
X_m(q) &= \left[ \binom{m+1}{2} q^{4m+2} + \binom{m}{2} q^{4m+1} \right] (q+1) \\
&\quad - (q+1) \left\{ q^{2m+2} \left[ \frac{1-q^{2m}}{(1-q^2)^2} - \frac{mq^{2m}}{1-q^2} \right] + q^{2m+3} \left[ \frac{1-q^{2m-2}}{(1-q^2)^2} - \frac{(m-1)q^{2m-2}}{1-q^2} \right] \right\};
\end{aligned}$$

but this is insufficient — we need to do four kinds of boundary effects! On one side of  $M_0$  and  $M_m$  (see Fig. 12) would be

$$Y_m(q) = (q+1)^2 \left\{ \binom{m}{2} q^{4m-1} - q^{2m+1} \left[ \frac{1-q^{2m-2}}{(1-q^2)^2} - \frac{(m-1)q^{2m-2}}{1-q^2} \right] \right\}.$$

Similarly, for  $Z_m(q)$  (from  $L_0$  to  $L_m$ ) and  $U_m(q)$  (between  $M_0$  and  $L_m$ ):

$$\begin{aligned}
Z_m(q) &= \binom{m+1}{2} q^{4m+1} + m(m-1)q^{4m} + \binom{m}{2} q^{4m-1} - q^{2m+1} \left[ \frac{1-q^{2m-1}}{(1-q)^2} - \frac{(2m-1)q^{2m-1}}{1-q} \right] \\
U_m(q) &= \binom{m}{2} q^{4m-1} + (m-1)^2 q^{4m-2} + \binom{m}{2} q^{4m-3} - q^{2m} \left[ \frac{1-q^{2m-2}}{(1-q)^2} - \frac{(2m-2)q^{2m-2}}{1-q} \right].
\end{aligned}$$




Figure 12: A different “Edge Effect”, as seen before and after the squaring




Figure 13: The two other “Edge Effects”, before and after the squaring

Finally, we can finish the computation for  $\mathcal{W}(R_{n,k})$ , when  $n > k$ :

$$\mathcal{W}(R_{n,k}; q) = \mathcal{W}(\text{Cb}_{2n+1,2k}; q) + \Delta^* \mathcal{W}(R_{n,k}) - 2 \left\{ \sum_{\ell=1}^{k-1} (k - \ell) [X_\ell(q) + Y_\ell(q) + Z_\ell(q) + U_\ell(q)] \right\}.$$

The last pair of braces can be evaluated by expanding the summand in  $\ell$ ,  
then sum over  $\ell = 1..k - 1$  to get:

$$\begin{aligned}
& \sum_{\ell=1}^{k-1} [X_{\ell}(q) + Y_{\ell}(q) + Z_{\ell}(q) + U_{\ell}(q)] \\
= & \sum_{\ell=1}^{k-1} \left\{ \ell^2 q^{4\ell} \left[ \frac{(1+q)^2 (1+q^2)^2}{2q^3} \right] - \ell q^{4\ell} \left[ \frac{(1+q)(1+q^2)(1-2q-2q^2-2q^3+q^4)}{2q^3(1-q)} \right] \right. \\
& \left. + q^{4\ell} \left[ \frac{(1+q)^2}{(1-q)^2} \right] - q^{2\ell} \left[ \frac{(1+q)^2}{(1-q)^2} \right] \right\} \\
= & \frac{k^2 q^{4k+1}}{2(1-q)^2} + \frac{k (q^{4k+5} + 2q^{4k+4} + 2q^{4k+3} + 2q^{4k+2} + q^{4k+1} - 2q^3)}{2(1-q)^2 (1-q^4)} \\
- & \frac{q^2 (1-q^{2k}) [q^{2k}(q^6 + q^5 + 2q^4 + 3q^3 + 2q^2 + q + 1) - (q^5 + q^4 + q^3 + q^2 + q)]}{(1-q)^2 (1-q^4)^2}
\end{aligned}$$

We have arrived at the Wiener polynomial of one of the hex carpets:

**Theorem 12** *When  $n > k$ , we have*

$$\begin{aligned}
\mathcal{W}(R_{n,k}; q) &= \frac{[2q(q^{2n+1} - 1) - (2n+1)(q^2 - 1)][2q(q^{2k} - 1) - 2k(q^2 - 1)]}{2(q-1)^4} - k(2n+1) \\
&- \frac{nq}{(1-q)^3} \left[ (1-q^{2k})^2 (1+q) - 2k(1-q)(1-q^{4k}) \right] \\
&- \frac{2q^{4k+2}}{(1-q)^2 (1-q^2)} + \frac{kq}{(1-q)^2} + \frac{q^{2k+1}(q+1)}{(1-q)^3} \\
&+ \frac{q^{4k+2}(2q^4 + q^3 + 2q^2 + q + 2) - (q + 2q^3 + 2q^4 + 2q^5 + q^7)}{(1+q)(1-q)^3(1+q^2)^2} \\
&+ \frac{k^2 q^{4k+1}}{2(1-q)^2} + \frac{k (q^{4k+5} + 2q^{4k+4} + 2q^{4k+3} + 2q^{4k+2} + q^{4k+1} - 2q^3)}{2(1-q)^2 (1-q^4)} \\
&- \frac{q^2 (1-q^{2k}) [q^{2k}(q^6 + q^5 + 2q^4 + 3q^3 + 2q^2 + q + 1) - (q^5 + q^4 + q^3 + q^2 + q)]}{(1-q)^2 (1-q^4)^2}
\end{aligned}$$

There is an “explicit formula” for each of the hex carpets — finding Wiener polynomials of  $S_{n,k}$  and  $P_{n,k}$  (as defined in [12]) and their variants is

a similar process except for fiddling with extra vertices to bring the squaring up to a more symmetric grid. Each result will take up half a page, however.

## 4 Summary and Discussion

Even in the new millennium, Wiener numbers and polynomials is still a topic where interesting problems abound.

1. Wiener Polynomials for branched or fused polycyclic chains, nor for that matter polycyclic rings has currently known non-recursive derivation. Wiener numbers of few other chemically interesting families of graphs with structure in more than 1 dimension has been determined other than the regular hex carpets as defined in [12].
2. It is also desirable to extend formulas for Wiener numbers to Wiener polynomials, but many results cannot be easily generalized. There are several nice propositions which does not yet have  $q$ -analogs, such as the elegant Eq. 14 and similar results on trees.
3. A formula for the Wiener number of a hex crown (see [12]) was proved by the present authors ([34]), but computing the Wiener polynomial is not easy due to the same problem that precluded Eq. 14 from having a nice  $q$ -analog, despite the high degree of symmetry.

The problems mentioned above in extending results from Wiener numbers to polynomials are exemplified by the following two propositions, aside

from the aforementioned Merris-McKay result:

**Proposition 13** *The following holds for a tree  $T$*

1. *For each vertex  $x \in T$  that has degree at least 3 (a “branch point”), let  $F(x)$  be the set of components of the forest  $T \setminus \{x\}$ , then Gutman showed that*

$$\mathcal{W}(T) = \binom{|T|+1}{3} - \sum_{\substack{x \in V(T) \\ \deg(x) \geq 3}} \sum_{\{T_1, T_2, T_3\} \in \binom{F(x)}{3}} |T_1| |T_2| |T_3| \quad (14)$$

2. *It was shown by Wiener himself that if for each edge  $e$  we call  $n_1(e)$  and  $n_2(e)$  the number of vertices to either side of the edge  $e$ , then*

$$\mathcal{W}(T) = \sum_{e \in T} n_1(e) n_2(e).$$

Eq. 14 has a very good combinatorial interpretation and useful applications. Most recently ([29]) it was used to identify the high alkane isomer with the smallest Wiener Index. but no analogous formula for Eq. 14 or either of the two other elegant results in Wiener polynomials exist so far! Many proofs about Wiener numbers depends on a counting argument which may simply be absent present when handling generating functions. This explains why formulas about Wiener numbers just don't generalize very easily at all to Wiener polynomials.




Figure 14: Crown of Order 6

In short, there are still interesting work remains to be done.

## Acknowledgements

The authors would like to dedicate this work to the memory of the late Prof. Gian-Carlo Rota, to whose kind encouragement we owe much in this paper.

## References

- [1] D. Bonchev, *Information theoretic indices for characterization of chemical structure*, Research Studies Press(Chichester), 1983.
- [2] F. Buckley and F. Harary, *Distances in Graphs*, Addison-Wesley (New York), 1990.
- [3] J.-C. Chen, I. Gutman and Y.-N. Yeh, *On the sum of all distances in graphs*, Tamkang J. of Mathematics 25(1993), 83–86.

- [4] R. C. Entringer, D. E. Jackson, D. A. Snyder, *Distance in graphs*. Czechoslovak Math. J. 26(101) (1976), 283–296.
- [5] I. Gutman and O. E. Polansky, *Mathematical concepts in organic chemistry*, Springer-Verlag (Berlin), 1986
- [6] I. Gutman, Y.-N. Yeh, S.-L. Lee, and J.-C. Chen, *Wiener Numbers of Dendrimers*, MATCH 30(1994) 103-115.
- [7] I. Gutman, S.-L. Lee, Y.-L. Luo and Y.-N. Yeh, *Recent results in the theory of the Wiener number*, Indian J. of Chem., 32A (1993), 651-661.
- [8] I. Gutman and Y.-N. Yeh, *On the sum of all distances in composite graphs*, Discrete Math. 135 (1994), no. 1-3, 359–365.
- [9] F. Harary, *Status and contrastatus*, Sociometry 22 (1959), 23-43.
- [10] H. Hosoya, *Topological index. A newly proposed quantity characterizing the topological nature of structural isomers of saturated hydrocarbons*, Bull. Chem. Soc. Jpn. 44(1971), 2332-2339.
- [11] H. Hosoya, *On some counting polynomials in Chemistry*, Discrete Appl. Math. 19 (1988), 239-254.
- [12] I-W. Huang, B.-Y. Yang, and Y.-N. Yeh, *Wiener Indices of Hex Carpets— from Hexagon Models to Square Grids*, SE Asia Bull. of Math. 20(1996), 81-102.
- [13] W.-C. Huang, B.-Y. Yang, and Y.-N. Yeh, *From Ternary Strings to Wiener indices of Benzenoid Chains*, Discrete Appl. Math. 73 (1997), 113–131.

- [14] R. Merris, *Laplacian Matrices of Graphs: a Survey*, Linear Algebra Appl. 197/198 (1994), 143-176.
- [15] Z. Mihalić, D. Veljan, D. Amić, S. Nikolić, D. Plavšić, and N. Trinajstić, *The Distance Matrix in Chemistry*, J. of Math. Chem. 11(1992), 223-258.
- [16] B. Mohar, *Eigenvalues, Diameter, and Mean Distance in graphs*, Graphs and Combinatorics 7 (1991), 53-64.
- [17] B. Mohar and T. Pisanski, *How to compute the Wiener index of a graph*, J. Math. Chem. 2 (1988) 267-277.
- [18] I. Motoc and A. Balaban, *Topological indices: intercorrelation, physical meaning, correlational ability*, Revue Roumaine de Chimie 26 (1981), 593-600.
- [19] J. Plesnik, *On the sum of all distances in a graph or digraph*, J. Graph Theory 8 (1984), 1-21
- [20] O. Polansky, M. Randić, and H. Hosoya, *Transfer Matrix Approach to the Wiener Numbers of Catacondensed Benzenoids*, MATCH 24(1989) 3-28.
- [21] D. H. Rouvray, *On the sum of distance matrix elements*, in Chemical Applications of Graph Theory (A. T. Balaban, ed.), Academic Press (New York), 1976.
- [22] D. H. Rouvray, *The role of the topological distance matrix in chemistry*, in "Mathematics and Computational Concepts in Chemistry" (N. Trinajstić, ed.), Horwood Publishers (Chichester) 1986, 295-306.
- [23] D. H. Rouvray, *The modelling of chemical phenomena using topological indexes*, J. Comput. Chem. 8(1987), 470-480.

- [24] B. E. Sagan, Y.-N. Yeh, and P. Zhang, *The Wiener Polynomial of a Graph*, to appear in Int'l J. of Quantum Chem.
- [25] L. Šoltés, *Transmission in graphs: a bound and vertex removing*, Math. Slovaca 41 (1991), 11-16
- [26] R. P. Stanley, *Log-concave and unimodal sequences in algebra, combinatorics, and geometry*, in "Graph Theory and Its Applications, East and West", Ann. NY Acad. Sci. 576(1989), 500-535.
- [27] H.-H. Teh and S.-C. Shee, *Algebraic Theory of Graphs*, Lee Kong Chian Inst. of Math. and Comp. Sci. Nanyang Univ. Singapore.
- [28] N. Trinajstić, *Chemical Graph Theory*, Vol. 2, CRC Press, Boca Raton (1983),
- [29] S.-C. Liu, L.-D. Tung, and Y.-N. Yeh, *The Trees with Minimum Wiener Number*, to appear in Intl. J. of Quantum Chem.
- [30] H. Wiener, *Structural determination of paraffin boiling points*, J. A. Chem. Soc. 69 (1947), 17-20.
- [31] B.-Y. Yang, and Y.-N. Yeh, *Zigging and Zagging in Pentachains*, Advances in Applied Mathematics 16(1995) 72-94.
- [32] B.-Y. Yang, and Y.-N. Yeh, *Chains of Motley Gems and their Wiener Indices*, in Proc. of *First International Tainan-Moscow Algebra Workshop (Tainan, 1994)*, 329–349, de Gruyter (Berlin), 1996, Y. Fong *et al* ed., De Gruyter (Berlin).

- [33] B.-Y. Yang, and Y.-N. Yeh, *About Wiener Numbers and Polynomials*, to appear in Proc. of *Second International Tainan-Moscow Algebra Workshop (Tainan, 1996)*, to be published by de Gruyter (Berlin), Y. Fong *et al* eds.
- [34] B.-Y. Yang, and Y.-N. Yeh, *Wiener Number of a Hex Crown*, submitted.